





CONTRIBUTED PAPER

A national assessment of waterbird hunting in coastal wetlands of Suriname, South America

David S. Mizrahi¹  | Arie L. Spaans² | Marianne J. Spaans-Scheen² |
Amelia R. Cox³  | Benoit Laliberté³ | Eduardo Gallo-Cajiao^{4,5}  |
Christian Roy³ 

¹New Jersey Audubon, Bernardsville, New Jersey, USA

²Foundation Friends of Nature Conservation Suriname, Arnhem, Netherlands

³Canadian Wildlife Service, Environment and Climate Change Canada, Gatineau, Quebec, Canada

⁴School of Marine and Environmental Affairs, University of Washington, Seattle, Washington, USA

⁵Department of Human Dimensions of Natural Resources Management, Colorado State University, Fort Collins, Colorado, USA

Correspondence

David S. Mizrahi, New Jersey Audubon, Bernardsville, NJ, USA.
Email: david.mizrahi@njudubon.org

Present address

Amelia R. Cox, Northwest Territories Cumulative Impact Monitoring Program, Government of the Northwest Territories, Yellowknife, Northwest Territories, Canada.

Funding information

National Fish and Wildlife Foundation, Grant/Award Number: F14AP00790; Vogelbescherming Nederland; Environment and Climate Change Canada; Van der Hucht de Beukelaar Stichting; Stichting Dierenrampenfonds; het Cultuurfonds

Abstract

The central northern coast of South America has extensive wetlands critical for waterbird conservation. While waterbird harvest occurs in the region, the impact on species population dynamics remains unclear. This study assesses waterbird hunting in the coastal wetlands of Suriname, addressing: (i) the extent of waterbird harvest, (ii) changes in harvest magnitude over time, and (iii) the motivations and methods used by hunters. We collected data via a national survey of licensed hunters in 2006 and 2016 using structured interviews. A Bayesian hierarchical model was used to analyze the data. We estimated harvest levels for 11 species and three groups (small herons, small shorebirds, large shorebirds). For most species, mean harvest per hunter significantly decreased from 2006 to 2016, except for blue-winged teal and migratory shorebirds. Most hunting was for non-commercial purposes (personal consumption and recreational). This is the first national assessment of waterbird hunting in Suriname. Harvest levels vary by species, and the sustainability of these levels remains uncertain. Managing hunting in Suriname requires addressing both legal and illegal hunting. Given Suriname's importance for waterbirds, particularly species like the scarlet ibis and migratory shorebirds, it should be a priority for conservation efforts.

KEYWORDS

harvest management, scarlet ibis, shorebirds, Western Atlantic Flyway, western hemisphere, wildlife management

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2026 His Majesty the King in Right of Canada and The Author(s). *Conservation Science and Practice* published by Wiley Periodicals LLC on behalf of Society for Conservation Biology. Reproduced with the permission of the Minister of Environment and Climate Change.

1 | INTRODUCTION

Wetlands are biologically productive ecosystems that have sustained humans globally. The use of wetlands has included the harvesting of plants and animals, such as fish and waterbirds, with multiple purposes (e.g., cultural, subsistence, recreational, commercial). However, following increased pressure on the environment during the second phase of the Anthropocene (i.e., the Great Acceleration), the exploitation of some species may have contributed to population declines, and in some cases, extinction (Benitez-Lopez et al., 2017; Cornford et al., 2023; Steffen et al., 2011). Additionally, for long-distance migratory species, such as birds, harvest at specific locales can be compounded over large spatial scales and become unsustainable (e.g., Gallo-Cajiao et al., 2020). Consequently, it is necessary to evaluate harvest levels, their impacts on population dynamics, and the social and economic factors that drive them, in order to inform effective conservation and management policies for wetland-dependent species. Against this backdrop, there is a strong need to assess waterbird harvest worldwide, but particularly tropical regions, where people's livelihoods often rely on natural resources, governance capacity for wildlife management may be weak, and harvest monitoring is relatively limited (Amano et al., 2018; Brotherton et al., 2020).

The central northern coastal region of South America includes extensive wetlands crucial for waterbird species conservation (Butler et al., 2001; Murray et al., 2019). These ecosystems comprise mangrove forests, brackish lagoons, freshwater swamps, and the most extensive mudflats of South America, a flat coastline, and is strongly influenced by the Guiana Current. Noteworthy, this region has a low human footprint with little intertidal wetland losses (Murray et al., 2022), so threats associated with land use change are generally not prevalent (Allan et al., 2023). These coastal wetlands serve as important non-breeding and stopover areas for migratory shorebirds (i.e., Aves: Scolopacidae, Charadriidae) along the Western Atlantic Flyway. Additionally, over 100 species of waterbirds, many of which are non-migratory, occur in these coastal wetlands, including ducks (Anatidae), flamingoes (Phoenicopteridae), and herons (Ardeidae). Along the central northern coast of South America, Suriname has the largest concentration of some waterbird species (Morrison & Ross, 1989; Spaans, 1978), including scarlet ibis (*Eudocimus ruber*), semipalmated sandpiper (*Calidris pusilla*), greater yellowlegs (*Tringa melanoleuca*), lesser yellowlegs (*Tringa flavipes*), and short-billed dowitcher (*Limnodromus griseus*). The importance of this region has been recognized through the designation of Important Bird Areas by BirdLife International, Ramsar wetlands of international importance, and sites of hemispheric importance under the

Western Hemisphere Shorebird Reserve Network (BirdLife International, 2026; Ramsar Convention Secretariat, 1997; Western Hemisphere Shorebird Reserve Network, 2024).

Waterbird population trajectories in this region are generally unknown with some exceptions. For instance, the central northern coast of South America is not surveyed as part of the International Waterbird Census, hindering long-term monitoring of populations (Wetlands International, 2026). This data gap has also impeded assessments of protected area effectiveness, which is essential for management (Wauchope et al., 2022). However, some species that occur in this region are declining across their entire range, such as the scarlet ibis, wood stork (*Mycteria americana*), and white-cheeked pintail (*Anas bahamensis*; BirdLife International, 2025). Similarly, several migratory shorebird species (e.g., semipalmated sandpiper, lesser yellowlegs, and short-billed dowitcher) have shown a significant decline of over three decades along the central northern coast of South America (Morrison et al., 2012), consistent with overall declines observed in North America (Smith et al., 2023).

Harvest, whether for commercial, recreational or subsistence purposes, can significantly contribute to mortality rates and influence population dynamics of some waterbird species. Hence, harvest assessments can play a crucial role in the development of sustainable harvest management frameworks and to inform conservation objectives (Cox et al., 2024; Koneff et al., 2017; Watts et al., 2015). Particularly, harvest assessments are important for shorebirds, as they typically have slow life history traits, such as small clutch sizes and only one breeding attempt per season, making them highly vulnerable to high harvest levels or even to seemingly low harvest levels when populations are declining (Watts et al., 2015). For migratory bird species, harvest assessments must also occur in multiple countries to account for the cumulative effects of harvest to be estimated. Recent studies from Guyana (Andres et al., 2022), French Guiana (Taylor, 2017), and Brazil (Bosi de Almeida et al., 2018) show widespread shorebird hunting in the central northern coast of South America. Beyond this specific region, migratory shorebirds have also been hunted along the Western Atlantic Flyway in parts of the Lesser Antilles (Gebauer, 2018; Reed et al., 2018; Watts & Turrin, 2016). In the United States and Canada, the extent of hunting remains unclear, as part of subsistence hunting (but see Naves et al., 2019) and illegal harvest during recreational hunting (Katzner et al., 2020).

In addition to evaluating harvest from an ecological standpoint, understanding the human dimensions of hunting is crucial for informing conservation policy. These dimensions include parameters such as demographics of resource users (e.g., age, gender, income), hunting's contribution to livelihoods, and attitudes

toward hunting and regulations (Eriksson et al., 2023; Vaske & Manfredi, 2012). For instance, hunting of migratory shorebirds in Southeast Asia can make a considerable contribution to household income (Gallo-Cajiao et al., 2020; Zöckler et al., 2010). In Guyana, shorebird hunters are typically fishermen supplementing their income when fishing conditions are poor (Andres et al., 2022). Understanding these parameters in the context of the central northern coast of South America could help inform more effective and equitable harvest management policies.

Here, we present a case study on a national assessment of waterbirds harvested in coastal wetlands of Suriname. Our aim is to contribute to the evaluation of harvest sustainability of waterbird species in Suriname within the context of central northern South America and the Western Atlantic Flyway. Specifically, we ask: (i) what is the recent extent of waterbird harvest in coastal wetlands of Suriname? (ii) has the magnitude of waterbird harvest changed over time? and (iii) what are the motivations of waterbird hunters and what harvest methods they use?

2 | METHODS

2.1 | Study area

Suriname, located on the central northern coast of South America, covers 163,800 km² and has a 386 km coastline.

It borders the Atlantic Ocean, Guyana, French Guiana, and Brazil. The country has a tropical climate with two dry and two wet seasons. Major rivers include the Courantyne, Coppename, Saramacca, Suriname, Commewijne, and Marowijne. Most of Suriname's population (87%) lives in the coastal area, where the economy is primarily driven by mining and agriculture (Food and Agriculture Organization, 2024). The coastal zone, mostly low and flat, comprises the lower and upper coastal plains, which are key waterbird habitats (Spaans et al., 2018). Wetland ecosystems in the lower coastal plain include mudflats, beaches, estuaries, mangroves, salt pans, coastal marshes, swamp forests, and shallow brackish lagoons. The upper coastal plain includes herbaceous and peat swamps. Our study focused on all coastal districts except Marowijne, where hunters were difficult to reach (Figure 1).

The coastal zone of Suriname has various designations under different international institutional frameworks, including one wetland of international importance under the Ramsar Convention, three hemispherically important sites under the Western Hemisphere Shorebird Reserve Network, and four Important Bird Areas under BirdLife International (BirdLife International, 2026; Ramsar Convention Secretariat, 1997; Western Hemisphere Shorebird Reserve Network, 2024). Domestically, there are four Multiple-Use Management Areas (MUMAS) and four nature reserves as well (Spaans et al., 2018). Some of these areas have overlapping designations.

FIGURE 1 Map of Suriname with district borders, showing coastal districts numbered: (1) Nickerie, (2) Coronie, (3) Saramacca, (4) Wanica, (5) Paramaribo, (6) Commewijne, and (7) Marowijne. The Marowijne district and non-coastal districts (Para [purple], Brokopondo [yellow] and Sipaliwini [green]) were not surveyed during this study.



TABLE 1 Species and species groups of waterbirds included in the hunter survey conducted in six districts of Suriname in 2006 and 2016.

Common name	Scientific name	IUCN listing	Hunting status according to Suriname's law	
			Migratory	
Blue-winged teal	<i>Spatula discors</i>	LC	Yes	Game species
White-cheeked pintail	<i>Anas bahamensis</i>	LC	No	Game species
Black-bellied whistling-duck	<i>Dendrocygna autumnalis</i>	LC	No	Game species
Muscovy duck	<i>Cairina moschata</i>	LC	No	Game species
American flamingo	<i>Phoenicopterus ruber</i>	LC	No	Protected
Cocoi heron	<i>Ardea cocoi</i>	LC	No	Protected
Great egret	<i>Ardea alba</i>	LC	No	Protected
Wood stork	<i>Mycteria americana</i>	LC	No	Protected
Jabiru	<i>Jabiru mycteria</i>	LC	No	Protected
Scarlet ibis	<i>Eudocimus ruber</i>	LC	No	Protected
Roseate spoonbill	<i>Platalea ajaja</i>	LC	No	Protected
Small herons (e.g., little blue heron, snowy egret, cattle egret, tricolored heron, black-crowned/yellow-crowned night-herons, boat-billed heron)	<i>Egretta caerulea</i> , <i>Egretta thula</i> , <i>Ardea ibis</i> , <i>Egretta tricolor</i> , <i>Nycticorax nycticorax</i> , <i>Nyctanassa violacea</i> , <i>Cochlearius cochlearius</i>	–	–	Protected
Small shorebirds (e.g., semipalmated and least sandpiper, semipalmated plover)	<i>Calidris pusilla</i> , <i>C. minutilla</i> , <i>Charadrius semipalmatus</i>	–	Yes	Protected
Large shorebirds (e.g., greater and lesser yellowlegs, short-billed dowitcher, willet, whimbrel, black-bellied plover)	<i>Tringa melanoleuca</i> , <i>T. flavipes</i> , <i>T. semipalmata</i> , <i>Limnodromus griseus</i> , <i>Numenius hudsonicus</i> , <i>Pluvialis squatarola</i>	–	Yes	Protected

Note: International Union for Conservation of Nature's (IUCN) Red List of Threatened Species information was sourced from the version 2022-2 (IUCN, 2022).

The Hunting Act 1954 and the corresponding implementing legislation (i.e., Hunting Decree 2002 as amended in 2009), determine the species of wildlife that may be hunted and when. All bird species occurring in Suriname are fully protected, except those designated as “game,” “cage birds,” and “predominantly harmful species.” Waterbird species without full protection under Suriname regulations and considered in this paper are: black-bellied whistling-duck (*Dendrocygna autumnalis*), muscovy duck (*Cairina moschata*), white-cheeked pintail, blue-winged teal (*Spatula discors*), Neotropic cormorant (*Nannopterum brasilianum*), and anhinga (*Anhinga anhinga*; Table 1).

2.2 | Study design and data collection

This is a cross-sectional and longitudinal single case study based on surveys conducted at two points in time that are 10 years apart. Travel is difficult, particularly on unpaved roads during the rainy season in Suriname, so data were collected in April 2006 and 2016, during what is typically the short dry season, to improve our chances of reaching hunters in their communities.

At the time the surveys were conducted, no ethics approval was required in Suriname. The surveys were conducted in collaboration with the Suriname Forest Service/Nature Conservation Division, a department within the Ministry of Land Policy and Forest Management. Staff from the agency were actively involved in survey implementation.

In 2016, potential participants were identified from hunter licensing records provided by the District Commissioners in each of the included districts. Neither the District Commissioners nor their staff were informed of the identities of the selected participants. No minors were involved in the study, as individuals under the age of 18 are not permitted to hold a hunting license in Suriname. The participant pool consisted of those who could be reached and agreed to participate. Due to challenges in contacting hunters, we could not create a stratified sample proportional to the number of hunters in each district.

Prior to administering the survey, all participants were informed of the study's purpose and how their responses would be used. They were advised that submission of a completed survey would be considered as

informed consent to participate. While participants were encouraged to answer all questions, they were also clearly informed that responding to any specific question was entirely voluntary. All surveys were administered in Dutch, the official language of Suriname. Survey responses were recorded on paper and later digitized. All responses remained anonymous; no names or identifying information were included on the questionnaires. Because we did not record personal identification information, we could not intentionally interview any hunters from the 2006 survey in 2016.

The survey questions in 2006 focused primarily on hunting frequency, magnitude and purpose, and species or group of species of waterbirds harvested (Supporting Methods 1). For some taxa, species reporting was not possible, or at least not accurately as hunters do not distinguish between species, so for these we present data for groups of species which share broad morphological and ecological traits (Table 1). The 2016 survey used all the same questions used in 2006 but included additional questions to improve our understanding of the hunter community (e.g., age, family members, hunting purpose), hunting seasonality, and hunting method (Supporting Methods 2).

Survey respondents self-reported their harvest, a common method for assessing waterbird hunting (Geissler, 1990; Smith et al., 2022). However, this approach has potential limitations, including recruitment bias and underreporting of mortality, such as wounded birds which may die later (i.e., crippling losses). In 2006, we piloted group surveys with hunters, but found that participants were conferring with each other, making the results unreliable. As a result, we switched to individual one-on-one interviews at hunters' homes, continuing this method in 2016. If interviewers suspected dishonesty, they cross-checked answers with game wardens familiar with the hunters' activities. Data deemed unreliable were excluded from analyses. All data were kept anonymous and confidential, and participation was voluntary.

2.3 | Data analysis

2.3.1 | Estimation of licensed hunters in 2016

In 2006, there was no reliable centralized hunter license database, so the total number of licensed hunters that year is unknown. For 2016, we used the Suriname hunter license database, which tracks only license registrations. According to Suriname law, hunting licenses are valid for up to 5 years, from January 1st to December 30th. We considered the total number of licensed hunters in 2016

as the unique registrations between January 1, 2011, and December 30, 2016. This method does not account for licenses that may have been suspended before 2016, as the database lacked records on renewals or suspensions, and no corrective measures were applied.

2.3.2 | Estimation of hunting levels

We used a Bayesian hierarchical model to analyze the harvest survey data (Supporting Methods 3). We modeled harvest data for 11 waterbird species and four taxa (i.e., small herons, small shorebirds, large shorebirds, and all shorebirds). Because of the limited number of respondents in each region, we pooled all the data together and did not consider variation in hunting across regions/districts. In 2006, survey respondents were asked to provide estimates for all shorebird species combined. In 2016, respondents were asked to distinguish between harvest of large and small shorebird species. Therefore, the survey of small and large shorebirds was set to zero for 2006 and we modeled the harvest of all shorebirds directly. For 2016, we instead modeled the harvest of small and large shorebirds and derived the harvest of all shorebirds from those two categories.

During the interview process, hunters reported the number of birds they harvested that year via 8 categories (either 0; 1; 2–10; 11–50; 51–100; 101–500; 501–1000; or >1000 birds). We therefore modeled the reported harvest of each species i by each individual hunter interviewed j via a categorical distribution:

$$\text{Harvest}_{i,j} \sim \text{Categorical}(\pi_{1:8,j,t}) \quad (1)$$

where $\pi_{1:8}$ represents the vector of probabilities that the hunter will report their harvest in that category for a given species j in year t .

We assumed that the underlying distribution of harvest of each species would follow a negative binomial distribution, which allows for over dispersion that we observed in harvest data. We parametrized the negative binomial distribution via the mean and variance (Lindén & Mäntyniemi, 2011) and modeled the mean harvest $(\mu_{j,t})$ for a given species j in year t such as:

$$\log(\mu_{j,t}) = \eta_t + \varepsilon_{j,t}, \quad (2)$$

$$\varepsilon_{j,t} \sim \text{Normal}(0, \sigma_t), \quad (3)$$

where η_t is the average harvest across all species on the log scale and a $\varepsilon_{j,t}$ is species specific derivation from this

mean in this given year. The species specific derivations are drawn from a normal distribution with year specific standard distribution. This hierarchical formulation allowed us to share information across species (i.e., “borrowing the strength” from the ensemble) but to also consider the change in harvest level across years. We then parametrized the variance ($\sigma_{j,t}^2$) of the model in function of the mean such as:

$$\sigma_{j,t}^2 = \mu_{j,t} + \theta \mu_{j,t}^2, \quad (4)$$

where θ is the overdispersion parameter for the quadratic mean–variance relationship of the classic negative binomial regression (Lindén & Mäntyniemi, 2011). We used moment matching to link estimates of the mean and variance to the species-specific and year specific probability ($p_{j,t} = \frac{\mu_{j,t}}{\sigma_{j,t}^2}$) and size parameter ($r_{j,t} = \frac{\mu_{j,t}^2}{(\sigma_{j,t}^2 - \mu_{j,t})}$) and then linked the probability of the harvest categories with the cumulative probability distribution (CDF; $F(k, p_{j,t}, r_{j,t})$) of the negative binomial distribution. The first category of harvest represents the probability that harvest for a given species each year is exactly zero and can be derived directly from the CDF:

$$\pi_{1j,t} = F(0, p_{j,t}, r_{j,t}). \quad (5)$$

The probability of the harvest for categories 2 to 7 can be derived by subtracting the CDF at the lower bound of the category from the CDF at the upper bound of the category. For example, the probability that harvest is between 2 to 10 birds (i.e., Category 3) for a given species in a given year can be estimated via:

$$\pi_{3j,t} = F(10, p_{j,t}, r_{j,t}) - F(2, p_{j,t}, r_{j,t}). \quad (6)$$

The last harvest category ($\pi_{8j,t}$; > 1000 birds) is the probability that the harvest is larger than the lower bound of the category:

$$\pi_{8j,t} = 1 - F(1000, p_{j,t}, r_{j,t}). \quad (7)$$

Hunters did not always complete the survey and did not report their harvest for all the species. In those cases, we used a data augmentation scheme and let the model assign a harvest category for the hunter. Because of the hierarchical formulation of the model, the category of harvest assigned to this hunter by the model will tend to

be close to the most frequent category of harvest observed for that species during that year.

Finally, we derived the total annual harvest in 2016 by multiplying the number of registered hunters with the mean annual harvest ($\mu_{j,2016}$) for a given species j . Since the Suriname hunter license database was incomplete for 2006, we were not able to estimate the number of registered hunters and total annual harvest in 2006.

We used R version 4.2.0 for data processing (R Core Team, 2020) and JAGS version 4.3.0 (Plummer, 2003), called via the R package jagsUI (Kellner, 2021) for analyses. We used non-informative priors for all parameters and ran three chains with randomized initial start values for 69,500 iterations, using the first 2000 iterations as burn in and thinning the chains to save every 12th iteration, with 1000 iterations of adaption yielding 45,000 posterior samples for each parameter. We assessed model convergence visually and by making sure that the Gelman–Rubin statistic (\hat{R}) was less or equal than 1.1 (Gelman et al., 2013). We also used a posterior predictive check to evaluate the performance of the model. All results are reported as the median with 90% Bayesian Credible Intervals, calculated as high-density intervals using *bayestestR* package (Makowski et al., 2019).

3 | RESULTS

3.1 | Hunters in Suriname

Surveys were conducted in six districts in 2006 and five districts in 2016, with the highest number of participants from Nickerie, Saramacca, and Commewijne (Table 2). Additionally, we were unable to recruit the same number of hunters from each district over the two time periods. Paramaribo District, despite having the largest population of licensed hunters, had the fewest participants (Table 2).

In 2006, 276 licensed hunters answered the survey (Table 2) but we excluded 45 of them from the analysis as

TABLE 2 Numbers of hunters interviewed about their harvest of waterbirds per district in Suriname in 2006 and 2016.

District	2006	2016
Nickerie	115	186
Coronie	8	21
Saramacca	100	29
Wanica	3	11
Paramaribo	3	0
Commewijne	47	42
Total	276	289




FIGURE 2 Mean birds harvested in a year (μ) per hunter based on surveys of hunters in Suriname in 2006 and 2016. Data points represent the median, with error bars indicating the 90% high-density intervals. American Ornithological Society species codes are presented in Table 3. Note that in 2006, hunters reported all shorebird species combined, but in 2016 large and small shorebirds were distinguished.

the answers were deemed to be unreliable. Of the remaining 231 surveys, only 69% ($n = 159$) of the hunters reported their harvest of all species included in the survey. Hunters completed the survey and reported their harvest most frequently for muscovy duck (91%), black-bellied whistling-duck (91%), white-cheeked pintail (90%), while response rates were lower for jabiru (*Jabiru*

mycteria; 82%) and American flamingo (*Phoenicopterus ruber*; 81%).

In 2016, a total of 289 male hunters answered the survey (Table 2) out of the 4727 hunters that were registered in the Suriname hunter license database. The answers to 16 surveys were deemed to be unreliable so we kept 273 surveys for the analysis. The ages of the respondents ranged from 19 to 85 years, with a mean age of 49 (SD = 11). Of the 273 surveys, 99% ($n = 270$) of the hunters reported their harvest for all species included in the survey. Of the surveyed hunters, 190 hunters indicated that they did not harvest any birds, 72 hunters reported harvesting 5 species or less, 8 hunters harvested between 6 and 10 species, and 3 hunters harvested more than 10 species.

In addition to the 11 species and four taxa that we modeled, hunters also reported the harvest of Neotropical cormorant, anhinga, brown pelican (*Pelecanus occidentalis*), black skimmer (*Rynchops niger*), and undefined Laridae taxa (i.e., *gulls and terns*). However, because the harvest levels for these species were too low for statistical modeling, we excluded them from the analysis.

3.2 | Estimated waterbirds harvest in Suriname

3.2.1 | Average harvest for 2006 and 2016

In both 2006 and 2016, mean harvest per hunter of all shorebirds was higher than for any other species or group (Figure 2). Results from the 2016 survey indicate that hunters harvested on average more small shorebirds than large shorebirds. After shorebirds, the most harvested species in 2006 were black-bellied whistling-duck, scarlet ibis, and cocoi heron; the jabiru and American flamingo were the least harvested species (Figure 2). In 2016, the most harvested species outside of shorebirds was blue-winged teal followed by black-bellied whistling-duck, while American flamingo and the jabiru were still the two least harvested species (Figure 2).

3.2.2 | Has hunting of waterbird species changed over time?

For most waterbird species, mean harvest per hunter declined between 2006 and 2016 (Figure 2). The exceptions were the blue-winged teal and the total migratory shorebirds (i.e., small and large combined), for which harvest did not decrease significantly between 2006 and 2016. In particular, wood stork, roseate spoonbill (*Platalea ajaja*), and jabiru harvest declined below one bird per

TABLE 3 Estimated total waterbird harvest by all registered hunters in Suriname in 2016.

Family/species	Species code	2016 annual harvest (90% BCI)
Anatidae		
Black-bellied whistling-duck (<i>Dendrocygna autumnalis</i>)	BBWD	31,824 (16955–50,389)
Blue-winged teal (<i>Spatula discors</i>)	BWTE	46,974 (24505–76,414)
Muscovy duck (<i>Cairina moschata</i>)	MUDU	6075 (3191–9713)
White-cheeked pintail (<i>Anas bahamensis</i>)	WCHP	18,520 (8987–30,855)
Phoenicopteridae		
American flamingo (<i>Phoenicopterus ruber</i>)	AMFL	512 (81–1505)
Ardeidae		
Cocoi heron (<i>Ardea cocoi</i>)	COHE	5129 (2115–9566)
Great egret (<i>Ardea alba</i>)	GREG	7631 (2470–16,363)
Small herons	SMHE	13,960 (3769–32,142)
Ciconiidae		
Jabiru (<i>Jabiru mycteria</i>)	JABI	92 (10–241)
Wood stork (<i>Mycteria americana</i>)	WOST	1230 (413–2408)
Threskiornithidae		
Roseate spoonbill (<i>Platalea ajaja</i>)	ROSP	865 (186–2262)
Scarlet ibis (<i>Eudocimus ruber</i>)	SCIB	19,124 (7667–36,053)
Charadriidae, Recurvirostridae, Scolopacidae		
Large shorebirds	Large	58,915 (23218–109,019)
Small shorebirds	Small	107,042 (36968–202,281)
Total shorebirds	Total	173,700 (93970–276,450)

Note: Estimates are presented as medians with 90% Bayesian Credible Intervals (BCI).

hunter on average in 2016. Hunters surveyed in 2016 generally agreed that their overall harvest had declined. Among respondents who answered this question and hunted in both 2006 and 2016, most reported hunting less in 2016 (77%), while a small minority (7%) reported hunting more than in 2006. When asked why they had changed their hunting practices, out of 66 responses provided

the most common response was that there were fewer birds available for harvest than there were in the past (29%). Some individuals also cited the increase in hunting regulations and increased enforcement (12%) and associated costs (6%).

3.2.3 | Total derived harvest 2016

Shorebirds were the most harvested group in general in 2016 with a total harvest of approximately 174 thousand birds (90% BCI = 94–276 thousand; Table 3). Small shorebirds (107 [90% BCI = 37–202] thousand) were harvested nearly twice as much as large shorebirds (59 [90% BCI = 23–109] thousand). Ducks were the next group of species with the most harvest, with the blue-winged teal having the highest total harvest (47 [90% BCI = 25–76] thousand), followed by herons, egrets and ibis, with the scarlet ibis (19 [90% BCI = 8–36] thousand; Table 3) having the highest total harvest. Finally, the wood stork, roseate spoonbill, American flamingo, and jabiru were less frequently harvested species with only the wood stork (1230 [90% BCI = 413–2408]; Table 3) harvest exceeding a thousand birds.

3.2.4 | What are the motivations of waterbird hunters and what hunting methods do they use?

Across both survey years (2006 and 2016), 63% of respondents traveled to their hunting location with the intention of engaging in another activity, generally fishing or recreation. Based on data from 2016 only, nearly all hunters who reported their hunting method ($N = 90$) reported hunting primarily with a firearm (99%), while a small proportion reported using a net (1%). In 2016, hunters who reported their primary purpose for hunting ($N = 90$) indicated that it was predominantly non-commercial in purpose, which includes personal consumption and recreational activities. Though there is overlap between these two motivations, the majority of hunters reported hunting for the former (70%) and only a small proportion for the latter (13%). Additionally, 13% of the hunters reported selling the birds, inferring a commercial purpose.

4 | DISCUSSION

Harvest of waterbird species is pervasive across coastal wetlands of Suriname, has declined over time, and is primarily practiced for personal consumption. To the best of

our knowledge, this is the first national assessment of waterbird harvest from Suriname, part of a region of critical importance to shorebird conservation in central northern South America and along the Western Atlantic Flyway. While the importance of waterbird harvest in Suriname has been internationally recognized for many years (AFSI Harvest Working Group, 2017, 2020; Ottema & Spaans, 2008), no structured assessment has been previously conducted. Waterbird harvest occurs along the entire coast of Suriname and includes species across a wide range of taxonomic groups. Among all waterbirds hunted in Suriname, the scarlet ibis and migratory shorebirds are of particular conservation concern due to higher hunting levels, declining populations, and slow life history traits (BirdLife International, 2016; Olmos & Silva e Silva, 2001; Smith et al., 2023; Watts et al., 2015).

In Suriname, waterbirds are primarily harvested using firearms, with only a small proportion using alternative methods, such as nets and choking wires. This pattern is similar to French Guiana (Taylor, 2017) but contrasts with neighboring Guyana and Brazil, where choking wires, fishing nets, and night lighting are the most common methods (Andres et al., 2022; Bos de Almeida et al., 2018). In line with broader understanding of wetland use by humans, waterbird harvest did not happen in isolation in our study but rather alongside other resource uses, particularly fishing.

Based on hunters' self-reported motivations, waterbird harvest in Suriname was primarily non-commercial and driven by recreational interest and personal consumption; the latter was the most frequently cited motivation in our study. This pattern contrasts with Guyana, where commercial harvest of migratory shorebirds occurs at a relatively larger scale although the overall harvest is less than in Suriname (Andres et al., 2022). While the limited commercialization reported in Suriname is encouraging, cross-border transport of waterbirds from western Suriname to markets in Guyana has been documented and warrants further investigation. Beyond subsistence, some of the harvest done for personal consumption may have cultural dimensions, as suggested by the consumption of birds such as the scarlet ibis at social celebrations (Ottema & Spaans, 2008), a practice noted elsewhere in the region as being driven by the bird's status as a traditional delicacy (BirdsCaribbean, 2017).

Species hunted represented a wide range of taxonomic groups and harvest levels varied between species, a pattern consistent with tropical wetlands elsewhere (Bos de Almeida et al., 2018; Deniau et al., 2022; Iskandar & Karlina, 2004; Ramachandran et al., 2017). Ducks were among the most heavily harvested groups, likely reflecting both their high availability and hunter

preferences for larger-bodied species. Of all the duck species hunted, only the blue-winged teal is considered migratory. Harvest levels of blue-winged teal from Suriname were modest compared to North America, representing about 1.68% of the combined harvest estimated for the US and Canada during the same year (Raftovich et al., 2018). The scarlet ibis was also among the species with the highest levels of hunting, and although it is hunted elsewhere in the tropics, the levels reported here are unusually high compared to recent reports from other regions (Deniau et al., 2022). Harvest of other species of waterbirds such as the American flamingo, the wood stork, the jabiru, and the roseate spoonbill was relatively low and in line with levels reported for other areas such as the Sahel (Deniau et al., 2022).

Current estimates of migratory shorebird harvest in Suriname exceed those available for any other country along the central northern coast of South America and the Lesser Antilles. We estimated that ≈ 174 thousand migratory shorebirds were harvested in 2016, but this is a conservative estimate as it does not account for crippling loss, under reporting because of illegality or unlicensed hunters. While empirical data on crippling rates for shorebirds, and particularly in South America, are virtually non-existent, estimates from other migratory game birds provide valuable context. For species such as doves and waterfowl, crippling losses typically range from 10% to over 25% and may be influenced by various hunting factors, including the type of ammunition used (i.e., steel shot vs. lead shot; Schulz et al., 2013; Ellis & Miller, 2022; Moreno-Zárate et al., 2023). Nevertheless, our estimate for shorebird harvest in Suriname greatly exceeds combined estimates reported for Guyana, the Lesser Antilles, and Brazil (AFSI Harvest Working Group, 2020; Andres et al., 2022; Bos de Almeida et al., 2018).

Our findings are limited by the qualitative, self-reported nature of the survey data, including recall bias, rounding, or intentional misreporting. Broad harvest categories and assumptions about the underlying distribution may affect precision but are unlikely to alter the overall magnitude of our estimates. Misclassification near range category boundaries and limited information on extreme values may affect estimation of the underlying distribution, and the assumption of a negative binomial distribution may not fully capture its shape, potentially biasing mean and total harvest estimates. We also assumed surveyed hunters were representative of all registered hunters. Future surveys could improve accuracy through narrower harvest categories and a stratified sampling.

The sustainability of waterbird harvest in Suriname remains largely unknown, but inferences can be drawn from ecological theory and species life-history traits. The

population-level effects of harvest range from compensatory to additive, or even super-additive mortality, depending on hunting intensity and species life history (Anderson & Burnham, 1976). Additive mortality occurs when harvest reduces survival, while compensatory mortality arises when harvest does not affect survival due to density dependence or targeting individuals that would have died naturally (Cooch et al., 2014; Sandercock et al., 2011). Generally, species with large clutch sizes, high natural mortality, and large populations experiencing density dependence can generally sustain higher harvest rates than species with small clutch sizes, high annual survival, and small populations (e.g., most shorebird species; Nichols et al., 1984; Cooch et al., 2014). The timing of harvest relative to natural mortality is also important, with hunting timed during or after periods of high natural mortality more likely to be additive (Blomberg, 2015; Kokko, 2001). For many species, migration represents a period of elevated natural mortality particularly for juveniles (Lok et al., 2015; Rushing et al., 2017; Sergio et al., 2019).

While duck hunting may pose relatively lower conservation concern, uncertainty around population status of resident birds in Suriname suggests a precautionary approach is warranted. The scarlet ibis population is declining across its range (Sainz-Borgo, 2025), and given its slow life history traits (i.e., small clutch size, delayed sexual maturity; Olmos & Silva e Silva, 2001), current harvest levels may be unsustainable. A more reliable assessment of the scarlet ibis population could encourage better adherence to regulations, as hunters may use uncertainty as a justification to dismiss the need for the law (i.e., neutralization theory; Eliason, 1999). Compliance promotion campaigns could help, but poachers often comply only in the presence of law enforcement (Jacoby, 2014; von Essen et al., 2014). Any harvest management strategies and compliance promotion efforts will need to consider broader socio-economic contexts such as local wetland use, livelihoods, and governance capacity to be effective (Ostrom, 2009; Sutherland et al., 2014).

For migratory shorebirds, species-level harvest estimates are lacking, but concern is particularly high for the semipalmated sandpiper, a Near Threatened species (BirdLife International, 2024) that likely accounts for a large percentage of the small shorebirds harvested in Suriname given its prevalence in the region (Morrison & Ross, 1989). Shorebird harvest in Suriname remains high, and up to 44% of the flyway-wide sustainable hunting levels could already be taken from neighboring Guyana (Andres et al., 2022; Watts et al., 2015).

Most species in our study are protected and the harvest of flamingoes, herons, storks, ibises, and shorebirds

in our surveys was illegal under Suriname's Hunting Act (1954). The observed declines for the harvest of flamingoes, herons, storks, and ibises are encouraging. These declines are also concurrent with an increase in compliance promotion and enforcement of harvest regulations, which likely contributed to the reduction in take. Shorebirds harvest level remains problematic. The status of shorebirds as protected species also complicates the engagement of hunters over species identification and creates challenges for the assessment of shorebird harvest sustainability (Solomon et al., 2015). This situation is not unique to Suriname; however, it has been observed elsewhere, such as in Guyana (Andres et al., 2022) and the East Asian-Australasian Flyway (Gallo-Cajiao et al., 2020).

Furthermore, a patchwork approach to regulating hunting of migratory shorebirds can blur its full effects until data are aggregated at a full life-cycle scale (Gallo-Cajiao et al., 2020). Regulatory frameworks for shorebird harvest in particular are variable across the north coast of South America (Watts & Turrin, 2016). Shorebirds are totally protected from harvest in Brazil, legally harvested for sport in French Guiana, and entirely unregulated in Guyana, including commercial hunting. This regulatory heterogeneity creates opportunities for laundering, whereby shorebirds illegally harvested in one jurisdiction may be sold in others where regulations are weak or absent. Although international agreements such as the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) and the Convention on Migratory Species (CMS) are relevant, their effectiveness is limited in this context. Suriname, for example, is a party to CITES but not CMS, while migratory shorebird species are listed under CMS but not CITES.

Addressing these challenges will require a coordinated, multi-jurisdictional approach grounded in improved, species-level harvest assessments and population monitoring, allowing harvest levels to be explicitly linked to population trends (Cox et al., 2025). Equally important is a better understanding of the human dimensions of shorebird harvest in the Western Atlantic Flyway, including equity, livelihood dependence, and perceptions of regulations and conservation measures, as policies lacking user acceptance are unlikely to succeed. Both lines of research on harvest effects and human dimensions will need to be further integrated into the policy process balancing conservation and human needs. More specifically, outputs from this research agenda could inform institutional arrangements already in place that have incorporated harvest mitigation as a priority, such as the Atlantic Flyway Shorebird Initiative (AFSI Harvest Working Group, 2017, 2020).

5 | CONCLUSION

Our study contributes to the understanding of wetland use more generally and to the consumptive use of waterbirds more specifically, both in the tropics as well as in Suriname within the context of central northern South America and the Western Atlantic Flyway. In doing so, we discovered that harvest could be happening at unsustainable levels both for resident and migratory species, supporting global analyses on the importance of overharvesting as a threat to biodiversity (Maxwell et al., 2016). Illegally hunted species are typically those that are of greatest conservation concern. The issue of waterbird harvest, and shorebirds in particular, has been recognized for decades in the central northern coast of South America and the Lesser Antilles (Hutt, 1991). However, structured and systematic assessments have only recently become available, highlighting the challenges involved in addressing this issue. More dedicated resources to monitor the status of populations and the level of harvest will greatly help develop more effective regulatory frameworks for waterbird conservation in the region. However, while enhancing existing regulatory frameworks is important, it is crucial to also prioritize efforts in promoting compliance, engaging with local communities, and developing alternative livelihoods. Given the importance of Suriname for waterbirds in the Americas, developing a cost-effective approach to Suriname's harvest management of species with international conservation concern, such as the scarlet ibis and migratory shorebirds, should remain a priority for the international community.

AUTHOR CONTRIBUTIONS

The conception of the project was led by David S. Mizrahi and Arie L. Spaans. The questionnaire was developed by Arie L. Spaans, Marianne J. Spaans-Scheen, and David S. Mizrahi. Data collection was led by Arie L. Spaans and Marianne J. Spaans-Scheen. Amelia R. Cox and Christian Roy analyzed the data and prepared the tables and figures. All the authors were involved in the interpretation of the data. The first draft of the paper was written by Eduardo Gallo-Cajiao and Amelia R. Cox. All authors approved the final version of the manuscript.

ACKNOWLEDGMENTS

We are deeply grateful to all the hunters who participated in this study. We also extend our sincere thanks to Marie Djosetro, Bryan Drakenstein, and Hesdy Esajas for their valuable contributions at the outset of the fieldwork in 2006 and 2016, respectively. Special thanks to Chanderbhan Dwarka, Amir Imraan, Astrid Moor, and Ashok Pherai for their assistance in conducting the surveys, Lena Usyk for compiling the survey results database and to Brendan Gray

for his support in compiling the hunter database. Fred Tremblay, two reviewers, and the editors also provided invaluable comments that greatly helped to improve this manuscript. The surveys conducted in 2006 received financial support from NVD Nature Conservation Fund, BirdLife Netherlands. Funding for the survey conducted in 2016 was provided to New Jersey Audubon by US Fish and Wildlife Service, through a grant from National Fish and Wildlife Foundation. The project received additional funding from het Cultuurfonds, Dierenrampenfonds Foundation, Van der Hucht De Beukelaar Foundation, and Environment and Climate Change Canada.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

ETHICS STATEMENT

Informed consent was obtained from all participants before interviews.

DATA AVAILABILITY STATEMENT

Data available upon request.

ORCID

David S. Mizrahi  <https://orcid.org/0000-0003-4338-2443>

Amelia R. Cox  <https://orcid.org/0000-0002-4804-5115>

Eduardo Gallo-Cajiao  <https://orcid.org/0000-0002-8607-2675>

Christian Roy  <https://orcid.org/0000-0002-5599-6234>

REFERENCES

- AFSI Harvest Working Group. (2017). *Achieving a sustainable shorebird harvest in the Caribbean and northern South America, progress report, 2011–2017* (Unpublished report). U.S. Fish & Wildlife Service, Migratory Bird Program. <https://www.shorebirdplan.org/wp-content/uploads/2018/08/AFSI-Shorebird-Hunting-Progress-Report-Sep-2017.pdf>
- AFSI Harvest Working Group. (2020). *Actions for the Atlantic flyway shorebird Initiative's shorebird harvest working group 2020–2025*. U.S. Fish & Wildlife Service, Migratory Bird Program. https://www.shorebirdplan.org/wp-content/uploads/2020/09/AFSI-Shorebird-Harvest-Action-Plan-2020_25-April-2020.pdf
- Allan, H., Levin, N., & Kark, S. (2023). Quantifying and mapping the human footprint across Earth's coastal areas. *Ocean & Coastal Management*, 236, 106476. <https://doi.org/10.1016/j.ocecoaman.2022.106476>
- Amano, T., Székely, T., Sandel, B., Nagy, S., Mundkur, T., Langendoen, T., Blanco, D., Soykan, C. U., & Sutherland, W. J. (2018). Successful conservation of global waterbird populations depends on effective governance. *Nature*, 553, 199–202. <https://doi.org/10.1038/nature25139>
- Anderson, D. R., & Burnham, K. P. (1976). *Population ecology of the mallard: VI. The effect of exploitation on survival* (Resource

- publication 128, Federal Government Series). U.S. Fish and Wildlife Service. <http://pubs.er.usgs.gov/publication/rp128>
- Andres, B. A., Moore, L., Cox, A. R., Frei, B., & Roy, C. (2022). A preliminary assessment of shorebird harvest in coastal Guyana. *Wader Study*, 129, 39–47. <https://doi.org/10.18194/ws.00262>
- Benitez-Lopez, A., Alkemade, R., Schipper, A., Ingram, D., Verweij, P., Eikelboom, J., & Huijbregts, M. (2017). The impact of hunting on tropical mammal and bird populations. *Science*, 356, 180–183. <https://doi.org/10.1126/science.aaj1891>
- BirdLife International. (2016). *Species factsheet: Scarlet ibis Eudocimus ruber*. <https://datazone.birdlife.org/species/factsheet/scarlet-ibis-eudocimus-ruber>
- BirdLife International. (2024). *Species factsheet: Semipalmated Sandpiper Calidris pusilla*. <https://datazone.birdlife.org/species/factsheet/semipalmated-sandpiper-calidris-pusilla>
- BirdLife International. (2025). *IUCN Red List for birds*. <https://datazone.birdlife.org>
- BirdLife International. (2026). *Country factsheet: Suriname*. <https://datazone.birdlife.org/country/factsheet/suriname>
- BirdsCaribbean. (2017). *Scarlet Ibis: A national symbol under siege*. <https://www.birdscaribbean.org/2017/08/scarlet-ibis-a-national-symbol-under-siege/>
- Blomberg, E. J. (2015). The influence of harvest timing on greater sage-grouse survival: A cautionary perspective. *The Journal of Wildlife Management*, 79, 695–703. <https://doi.org/10.1002/jwmg.877>
- Bosi de Almeida, J., Gallon de Aguiar, A., Damasceno, J. P., & Carvalho, M. R. (2018). *Assessment of shorebird hunting in northern Brazil* (Unpublished report). Sociedade para a Conservação das Aves do Brasil.
- Brotherton, S., Joyce, C. B., & Scharlemann, J. P. (2020). Global off-take of wild animals from wetlands: Critical issues for fish and birds. *Hydrobiologia*, 847, 1631–1649. <https://doi.org/10.1007/s10750-020-04222-2>
- Butler, R. W., Davidson, N. C., & Morrison, R. G. (2001). Global-scale shorebird distribution in relation to productivity of near-shore ocean waters. *Waterbirds*, 24, 224–232. <https://doi.org/10.2307/1522030>
- Cooch, E. G., Guillemain, M., Boomer, G. S., Lebreton, J.-D., & Nichols, J. D. (2014). The effects of harvest on waterfowl populations. *Wildfowl, Special Issue*, 4, 220–276.
- Cornford, R., Spooner, F., McRae, L., Purvis, A., & Freeman, R. (2023). Ongoing over-exploitation and delayed responses to environmental change highlight the urgency for action to promote vertebrate recoveries by 2030. *Proceedings of the Royal Society B: Biological Sciences*, 290, 20230464. <https://doi.org/10.1098/rspb.2023.0464>
- Cox, A. R., Gallo-Cajiao, E., Tremblay, F., Rateau, F., Uroy, K., Laliberté, B., Euphrosine, D., Lavanne, T., & Roy, C. (2025). *Legal harvest of shorebirds and resident game birds on a Caribbean Island: A Martinique case study* (Preprint). bioRxiv <https://doi.org/10.1101/2025.06.17.660260>
- Cox, A. R., Roy, C., Hanson, A., & Robertson, G. J. (2024). Canadian murre harvest management in the face of uncertainty: A potential biological removal approach. *The Journal of Wildlife Management*, 88, e22573. <https://doi.org/10.1002/jwmg.22573>
- Deniau, C. P., Mathevet, R., Gautier, D., Besnard, A., Cornu, G., & Le Bel, S. (2022). Flyways to hell? An empirical assessment of Palearctic migratory waterbird harvest practices in key wetlands of Sahel-sub-Saharan Africa. *Biological Conservation*, 265, 109,419. <https://doi.org/10.1016/j.biocon.2021.109419>
- Eliason, S. L. (1999). The illegal taking of wildlife: Toward a theoretical understanding of poaching. *Human Dimensions of Wildlife*, 4, 27–39.
- Ellis, M. B., & Miller, C. A. (2022). The effect of a ban on the use of lead ammunition for waterfowl hunting on duck and goose crippling rates in Illinois. *Wildlife Biology*, 2022(2), e01001. <https://doi.org/10.1002/wlb3.01001>
- Eriksson, L., Månsson, J., Liljebäck, N., Sandström, C., Johansson, M., Eklund, A., & Elmberg, J. (2023). The importance of structural, situational, and psychological factors for involving hunters in the adaptive flyway management of geese. *Scientific Reports*, 13, 7112. <https://doi.org/10.1038/s41598-023-34316-x>
- Food and Agriculture Organization of the United Nations. (2024). *Country profile—Suriname: Economy, agriculture and food security*. FAO Knowledge Repository. <https://openknowledge.fao.org/handle/20.500.14283/ca0427en>
- Gallo-Cajiao, E., Morrison, T. H., Woodworth, B. K., Lees, A. C., Szabo, J. K., & Fuller, R. A. (2020). Extent and potential impact of hunting on migratory shorebirds in the Asia-Pacific. *Biological Conservation*, 246, 108582. <https://doi.org/10.1016/j.biocon.2020.108582>
- Gebauer, M. (2018). *Historic and current shorebird hunting and management structure in Trinidad and Tobago* (Unpublished report). Environment and Climate Change Canada, Canadian Wildlife Service.
- Geissler, P. (1990). Estimation of confidence intervals for federal waterfowl harvest surveys. *Journal of Wildlife Management*, 54, 201–205.
- Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., & Rubin, D. B. (2013). *Bayesian data analysis* (3rd ed.). CRC Press/Chapman & Hall.
- Hutt, M. B. (1991). Shooting of migrating shorebirds in Barbados. In T. Salathé (Ed.), *Conserving migratory birds* (pp. 77–91). International Council for Bird Preservation.
- International Union for Conservation of Nature (IUCN). (2022). *The IUCN Red List of Threatened Species*. Version 2022–2. <https://www.iucnredlist.org/>
- Iskandar, S., & Karlina, E. (2004). Kajian pemanfaatan jenis burung air di pantai utara Indramayu, Jawa Barat. *Buletin Plasma Nut-fah*, 10, 43–48.
- Jacoby, K. (2014). *Crimes against nature: Squatters, poachers, thieves, and the hidden history of American conservation*. University of California Press.
- Katzner, T., Nelson, D. M., Diffendorfer, J. E., Duerr, A. E., Campbell, S. P., Leslie, D., & Yee, J. L. (2020). Illegal killing of nongame wildlife and recreational shooting in conservation areas. *Conservation Science and Practice*, 2, e279. <https://doi.org/10.1111/csp2.279>
- Kellner, K. (2021). *jagsUI: A wrapper around “rjags” to streamline “JAGS” analyses* [R package]. <https://CRAN.R-project.org/package=jagsUI>
- Kokko, H. (2001). Optimal and suboptimal use of compensatory responses to harvesting: Timing of hunting as an example. *Wildlife Biology*, 7, 141–150. <https://doi.org/10.2981/wlb.2001.018>
- Koneff, M. D., Zimmerman, G. S., Dwyer, C. P., Fleming, K. K., Padding, P. I., Devers, P. K., Johnson, F. A., Runge, M. C., & Roberts, A. J. (2017). Evaluation of harvest and information

- needs for North American sea ducks. *PLoS One*, *12*, e0175411. <https://doi.org/10.1371/journal.pone.0175411>
- Lindén, A., & Mäntyniemi, S. (2011). Using the negative binomial distribution to model overdispersion in ecological count data. *Ecology*, *92*, 414–421. <https://doi.org/10.1890/10-1081.1>
- Lok, T., Overdijk, O., & Piersma, T. (2015). The cost of migration: Spoonbills suffer higher mortality during trans-Saharan spring migrations only. *Biology Letters*, *11*, 20140944. <https://doi.org/10.1098/rsbl.2014.0944>
- Makowski, D., Ben-Shachar, M. S., & Lüdtke, D. (2019). bayestestR: Describing effects and their uncertainty, existence and significance within the Bayesian framework. *Journal of Open Source Software*, *4*, 1541. <https://doi.org/10.21105/joss.01541>
- Maxwell, S. L., Fuller, R. A., Brooks, T. M., & Watson, J. E. (2016). Biodiversity: The ravages of guns, nets and bulldozers. *Nature*, *536*, 143–145. <https://doi.org/10.1038/536143a>
- Moreno-Zárate, L., Arroyo, B., Sardà-Palomera, F., Rocha, G., Bota, G., & Peach, W. J. (2023). Age ratio, crippling losses and factors affecting daily hunting bags of European turtle-dove in Spain: Implications for sustainable harvest management of a declining migratory species. *Science of the Total Environment*, *868*, 161192. <https://doi.org/10.1016/j.scitotenv.2022.161192>
- Morrison, R. G., Mizrahi, D. S., Ross, R. K., Ottema, O. H., de Pracontal, N., & Narine, A. (2012). Dramatic declines of Semipalmated Sandpipers on their major wintering areas in the Guianas, northern South America. *Waterbirds*, *35*, 120–134. <https://doi.org/10.1675/063.035.0112>
- Morrison, R. G., & Ross, R. K. (1989). *Atlas of Nearctic shorebirds on the coast of South America*. Canadian Wildlife Service Special Publication.
- Murray, N. J., Phinn, S. R., DeWitt, M., Ferrari, R., Johnston, R., Lyons, M. B., Clinton, N., Thau, D., & Fuller, R. A. (2019). The global distribution and trajectory of tidal flats. *Nature*, *565*, 222–225. <https://doi.org/10.1038/s41586-018-0805-8>
- Murray, N. J., Worthington, T. A., Fuller, R. A., Bi, J. J., Choi, C. Y., Clinton, N., & Lyons, M. B. (2022). High-resolution mapping of losses and gains of Earth's tidal wetlands. *Science*, *376*, 744–749. <https://doi.org/10.1126/science.abm9583>
- Naves, L. C., Keating, J. M., Tibbitts, T. L., & Ruthrauff, D. R. (2019). Shorebird subsistence harvest and indigenous knowledge in Alaska: Informing harvest management and engaging users in shorebird conservation. *The Condor*, *121*, duz023. <https://doi.org/10.1093/condor/duz023>
- Nichols, J., Conroy, M., Anderson, D., & Burnham, K. (1984). Compensatory mortality in waterfowl populations: A review of the evidence and implications for research and management. *Transactions of the North American Wildlife and Natural Resources Conference*, *49*, 535–554.
- Olmos, F., & Silva e Silva, R. (2001). Breeding biology and nest characteristics of the Scarlet Ibis in southeastern Brazil. *Waterbirds*, *24*, 58–67. <https://doi.org/10.2307/1522244>
- Ostrom, E. (2009). A general framework for analyzing sustainability of social-ecological systems. *Science*, *325*(5939), 419–422.
- Ottema, O. H., & Spaans, A. L. (2008). Challenges and advances in shorebird conservation in the Guianas, with a focus on Suriname. *Ornitología Neotropical*, *19*, 339–346.
- Plummer, M. (2003). JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. In *3rd International Workshop on Distributed Statistical Computing*.
- R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Raftovich, R., Chandler, S., & Fleming, K. (2018). *Migratory bird hunting activity and harvest during the 2016–17 and 2017–18 hunting seasons*. U.S. Fish and Wildlife Service.
- Ramachandran, R., Kumar, A., Gopi Sundar, K. S., & Bhalla, R. S. (2017). Hunting or habitat? Drivers of waterbird abundance and community structure in agricultural wetlands of southern India. *Ambio*, *46*, 613–620. <https://doi.org/10.1007/s13280-016-0861-4>
- Ramsar Convention Secretariat. (1997). *Information sheet on Ramsar wetlands (RIS): Coppenamemonding Nature Reserve*. Ramsar Sites Information Service. <https://rsis.ramsar.org/ris/304>
- Reed, E. T., Kardynal, K. J., Horrocks, J. A., & Hobson, K. A. (2018). Shorebird hunting in Barbados: Using stable isotopes to link the harvest at a migratory stopover site with sources of production. *The Condor*, *120*, 357–370. <https://doi.org/10.1650/CONDOR-17-134.1>
- Rushing, C. S., Hostetler, J. A., Sillett, T. S., Marra, P. P., Rotenberg, J. A., & Ryder, T. B. (2017). Spatial and temporal drivers of avian population dynamics across the annual cycle. *Ecology*, *98*, 2837–2850. <https://doi.org/10.1002/ecy.1983>
- Sainz-Borgo, C. (2025). Scarlet Ibis (*Eudocimus ruber*), version 2.0. In N. D. Sly (Ed.), *Birds of the world*. Cornell Lab of Ornithology. <https://doi.org/10.2173/bow.scaibi.02>
- Sandercock, B. K., Nilsen, E. B., Brøseth, H., & Pedersen, H. C. (2011). Is hunting mortality additive or compensatory to natural mortality? Effects of experimental harvest on the survival and cause-specific mortality of willow ptarmigan. *Journal of Animal Ecology*, *80*, 244–258. <https://doi.org/10.1111/j.1365-2656.2010.01769.x>
- Schulz, J. H., Bonnot, T. W., Millsbaugh, J. J., & Mong, T. W. (2013). Harvest and crippling rates of mourning doves in Missouri. *Wildlife Society Bulletin*, *37*, 287–292. <https://doi.org/10.1002/wsb.249>
- Sergio, F., Tavecchia, G., Tanferna, A., Blas, J., Blanco, G., & Hiraldo, F. (2019). When and where mortality occurs throughout the annual cycle changes with age in a migratory bird: Individual vs. population implications. *Scientific Reports*, *9*, 17352. <https://doi.org/10.1038/s41598-019-53531-z>
- Smith, A. C., Villeneuve, T., & Gendron, M. (2022). Hierarchical Bayesian integrated model for estimating migratory bird harvest in Canada. *The Journal of Wildlife Management*, *86*, e22160. <https://doi.org/10.1002/jwmg.22160>
- Smith, P. A., Smith, A. C., Andres, B., Francis, C. M., Harrington, B., Friis, C., Morrison, R. G., Paquet, J., Winn, B., & Brown, S. (2023). Accelerating declines of North America's shorebirds signal the need for urgent conservation action. *Ornithological Applications*, *125*, duad003. <https://doi.org/10.1093/ornithapp/duad003>
- Solomon, J. N., Gavin, M. C., & Gore, M. L. (2015). Detecting and understanding non-compliance with conservation rules. *Biological Conservation*, *189*, 1–4. <https://doi.org/10.1016/j.biocon.2015.04.029>
- Spaans, A. L. (1978). Status and numerical fluctuations of some North American waders along the Surinam coast. *Wilson Bulletin*, *90*, 60–83.
- Spaans, A. L., Ottema, O. H., & Ribot, J. H. J. M. (2018). *Field guide to the birds of Suriname* (2nd ed.). Koninklijke Brill.

- Steffen, W., Grinevald, J., Crutzen, P., & McNeill, J. (2011). The Anthropocene: From global change to planetary stewardship. *Ambio*, 40, 739–761. <https://doi.org/10.1007/s13280-011-0185-x>
- Sutherland, W. J., Gardner, T. A., Haider, L. J., & Dicks, L. V. (2014). How can local and traditional knowledge be effectively incorporated into international assessments? *Oryx*, 48(1), 1–2.
- Taylor, A. R. (2017). *Hunter surveys in the Mana rice fields, French Guiana* (Final report). Department of Geography and Environmental Studies, University of Alaska Anchorage.
- Vaske, J. J., & Manfredi, M. J. (2012). Social psychological considerations in wildlife management. In D. J. Decker, S. J. Riley, & W. F. Siemer (Eds.), *Human dimensions of wildlife management* (pp. 43–57). JHU Press.
- von Essen, E., Hansen, H. P., Nordström Källström, H., Peterson, M. N., & Peterson, T. R. (2014). Deconstructing the poaching phenomenon: A review of typologies for understanding illegal hunting. *British Journal of Criminology*, 54, 632–651. <https://doi.org/10.1093/bjc/azu022>
- Watts, B. D., Reed, E. T., & Turrin, C. (2015). Estimating sustainable mortality limits for shorebirds using the Western Atlantic Flyway. *Wader Study*, 122, 37–53. <https://doi.org/10.18194/ws.00002>
- Watts, B. D., & Turrin, C. (2016). Assessing hunting policies for migratory shorebirds throughout the Western Hemisphere. *Wader Study*, 123, 6–15. <https://doi.org/10.18194/ws.00023>
- Wauchope, H. S., Jones, J. P., Geldmann, J., Simmons, B. I., Amano, T., Blanco, D. E., Fuller, R. A., Johnston, A., Langendoen, T., Mundkur, T., & Nagy, S. (2022). Protected areas have a mixed impact on waterbirds, but management

helps. *Nature*, 605, 103–107. <https://doi.org/10.1038/s41586-022-04617-0>

Western Hemisphere Shorebird Reserve Network. (2024). *WHSRN site profile: Suriname coastal wetlands*. <https://whsrn.org/whsrn-sites/>

Wetlands International. (2026). *Waterbird Populations Portal*. <https://wpp.wetlands.org>

Zöckler, C., Hla, T. H., Clark, N., Syroechkovskiy, E., Yakushev, N., Daengphayon, S., & Robinson, R. (2010). Hunting in Myanmar is probably the main cause of the decline of the Spoon-billed Sandpiper *Calidris pygmeus*. *Wader Study Group Bulletin*, 117, 1–8.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Mizrahi, D. S., Spaans, A. L., Spaans-Scheen, M. J., Cox, A. R., Laliberté, B., Gallo-Cajiao, E., & Roy, C. (2026). A national assessment of waterbird hunting in coastal wetlands of Suriname, South America. *Conservation Science and Practice*, e70273. <https://doi.org/10.1111/csp2.70273>